

Exhibit A

Cellular Immune Responses to the Hepatitis B Virus Polymerase^{1,2}

Eishiro Mizukoshi,^{3*} John Sidney,[†] Brian Livingston,[‡] Marc Ghany,^{*} Jay H. Hoofnagle,^{*} Alessandro Sette,[†] and Barbara Rehermann^{4*}

CD4⁺ T cells play an important role in hepatitis B virus (HBV) infection by secretion of Th1 cytokines that down-regulate HBV replication, and by promoting CD8⁺ T cell and B cell responses. We have identified and characterized 10 CD4⁺ T cell epitopes within polymerase and used them to analyze the immunological effects of long-term antiviral therapy as compared with spontaneous recovery from HBV infection. Candidate epitopes were tested for binding to 14 HLA-DR molecules and in IFN- γ ELISPOT and cytotoxicity assays using peripheral blood lymphocytes from 66 HBV-infected patients and 16 uninfected controls. All 10 epitopes bound with high affinity to the most prevalent HLA-DR Ags, were conserved among HBV genomes, and induced IFN- γ responses from HBV-specific CD4⁺ T cells. Several epitopes contained nested MHC class I motifs and stimulated HBV-specific IFN- γ production and cytotoxicity of CD8⁺ T cells. HBV polymerase-specific responses were more frequent during acute, self-limited hepatitis and after recovery (12 of 18; 67%) than during chronic hepatitis (16 of 48 (33%); $p = 0.02$). Antiviral therapy of chronic patients restored HBV polymerase and core-specific T cell responses during the first year of treatment, but thereafter, responses decreased and, after 3 years, were no more frequent than in untreated patients. Decreased T cell responsiveness during prolonged therapy was associated with increased prevalence of lamivudine-resistant HBV mutants and increased HBV titers. The data provide a rationale for the combination of antiviral and immunostimulatory therapy. These newly described HBV polymerase epitopes could be a valuable component of a therapeutic vaccine for a large and ethnically diverse patient population. *The Journal of Immunology*, 2004, 173: 5863–5871.

The hepatitis B virus (HBV)⁵ is a noncytopathic DNA virus that causes both acute and chronic liver disease. Despite the availability of an effective vaccine for more than two decades, hepatitis B remains one of the 10 most common causes of death worldwide. More than 5% of the world population, i.e., 400 million people, are currently infected with HBV, and >250,000 people die each year from HBV-related liver cirrhosis and hepatocellular carcinoma (1). The development of antiviral and immunostimulatory therapies for HBV-infected patients therefore remains an important priority.

Recovery from acute hepatitis B is the result of a combination of cellular and humoral immune responses. Whereas neutralizing Abs against hepatitis B (HB) virus surface (s) Ag (HBsAg) appear after recovery, cellular immune responses are generally detectable before the synthesis of neutralizing Abs (2) and precede the rise of

serum alanine aminotransferase (ALT) levels (3) as well as clearance of HBeAg and HBsAg. They are mediated by cells of the innate immune response (4, 5) and by CD4⁺ and CD8⁺ T cells that clear HBV by cytolytic (6) and noncytolytic, cytokine-mediated (4, 7) mechanisms. After recovery, HBV-specific T cells persist in the blood for decades (8). In contrast, in chronic HBV infection, HBV-specific cellular immune responses are typically weak, narrowly focused, and rarely detectable in the peripheral blood (2, 9–11). However, they are not completely absent, because transient increases in the cellular immune response have been shown to precede increases of ALT activity and can be followed by the development of neutralizing Abs and spontaneous recovery (12). Therefore, it has been suggested that therapeutic induction of HBV-specific cellular immune responses may lead to recovery from chronic hepatitis B, and a lipopeptide-based vaccine with an HLA-A2-restricted HBV CD8⁺ T cell epitope and a tetanus toxoid CD4⁺ T cell epitope has been evaluated as experimental vaccine (13, 14). Although CD8⁺ T cell responses against the HBcore peptide could be induced in healthy, uninfected controls (15) as well as in patients with chronic hepatitis B (16), this was not sufficient to mediate viral clearance.

Because HBV-specific CD4⁺ T cells contribute to induction (17) and maintenance of Ag-specific CD8⁺ T cells, license dendritic cells to activate CD8⁺ effector T cells (17, 18), and provide help for activation and differentiation of B cells, the induction of HBV-specific CD4⁺ T cells is regarded as an important component of any immunomodulatory therapy. Whereas CD4⁺ T cell epitopes have been identified in the HBcore, HBs, and HBe Ag, the largest protein, HBV polymerase, has not been studied. This is a significant omission because HBV polymerase is a highly immunogenic CD8⁺ T cell target in acute self-limited hepatitis, even though it is produced in significantly smaller quantities than the other HBV Ags (2). Because polymerase is essential for the earliest steps in the

*Liver Diseases Section, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Department of Health and Human Services, Bethesda, MD 20892; [†]La Jolla Institute for Allergy and Immunology, and [‡]Epimmune, Inc., San Diego, CA 92121

Received for publication August 19, 2003. Accepted for publication August 20, 2004.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was funded in part by National Institutes of Health-National Institute of Allergy and Infectious Diseases Contract N01-AI-95362.

² This is Publication Number 541 from the La Jolla Institute for Allergy and Immunology.

³ Current address: First Department of Internal Medicine, Kanazawa University School of Medicine, Kanazawa, Ishikawa 920-8641, Japan.

⁴ Address correspondence and reprint requests to Dr. Barbara Rehermann, Liver Diseases Section, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, 10 Center Drive, Room 9B16, Bethesda, MD 20892. E-mail address: Rehermann@nih.gov

⁵ Abbreviations used in this paper: HBV, hepatitis B virus; HCV, hepatitis C virus; HB, hepatitis B; c, core; s, surface; ALT, alanine aminotransferase.

HBV life cycle, recognition of this Ag may limit early HBV spread, and its high degree of conservation may prevent viral escape via mutations in T cell epitopes. Finally, only polymerase-encoding plasmids and not envelope-encoding plasmids were able to break tolerance and to induce specific CD8⁺ T cells upon immunization of HBV transgenic mice (19). The characterization of CD4⁺ T cell responses against HBV polymerase, which is performed in the current study, does therefore fill an important gap in our understanding of the HBV-specific immune response and is relevant for the development of multi-epitope, immunostimulatory vaccines.

A second issue that is addressed by the current study is the effect of antiviral therapy on T cell responsiveness. Whereas some studies reported a restoration of HBV-specific T cell responses when persistently infected patients were treated with nucleoside analogs that inhibit HBV replication (20, 21), these results have not been confirmed in other studies (22), and most treated patients do not maintain viral control (23, 24). Thus, the hypothesis that high levels of HBV may suppress HBV-specific T cell responses is still controversial. Studying 66 patients with past or present HBV infection, we show that HBV polymerase-specific responses correlated with clinical and serological recovery from hepatitis B and with the duration and outcome of antiviral therapy. Thus, monitoring HBV polymerase-specific CD4⁺ T cells is of significant value in a diagnostic sense and in immunopathology studies investigating host-virus interaction. In addition, the data provide a rationale for the combination of antiviral and immunostimulatory therapy, and the described HBV polymerase epitopes could be a valuable component of such therapy for a large and ethnically diverse patient population.

Materials and Methods

Patient population

Sixty-six adult patients with past or present HBV infection were grouped as follows: group 1, 1 patient with acute, self-limited hepatitis B, who was positive for HBsAg and later seroconverted to anti-HBe and anti-HBs; group 2, 17 recovered patients who were negative for HBsAg, but reactive for anti-HB core (c)⁺anti-HBs⁺; group 3, 14 chronically infected patients who were positive for HBsAg and who were not being treated with lamivudine; and group 4, 34 chronically infected patients who were positive for HBsAg and who were receiving lamivudine treatment. Groups 3 and 4 were subdivided into patients with (groups 3a and 4a, respectively) or without HBeAg in serum (groups 3b and 4b, respectively). No patient had clinically decompensated cirrhosis. All subjects had been followed in the Liver Diseases Section, NIDDK, National Institutes of Health for >2 years, tested negative for Abs to HIV and hepatitis C virus (HCV), and gave written informed consent to this institutional review board-approved study. Preliminary results of the outcome of lamivudine therapy (100 mg daily) in a subgroup of these patients have been reported (25). Sixteen

healthy blood donors without any history of hepatitis and without HBsAg or anti-HBe in serum served as controls.

Laboratory and virologic testing

Serum HBsAg, HBeAg, and anti-HBe were detected with commercial immunoassays (Abbott Laboratories, Abbott Park, IL). Molecular typing of HLA DR alleles was performed on genomic DNA using standard site-specific oligonucleotide PCR. Serum HBV DNA was quantitated by branched DNA signal amplification assay (Chiron Corporation, Emeryville, CA) with a lower limit of detection of 0.7×10^6 genome equivalents (10^6 genome equivalents = 1 mEq) per milliliter (26). Samples with HBV DNA of <1 mEq/ml were tested by quantitative PCR (National Genetics Institute, Los Angeles, CA) with a lower limit of detection of 100 copies/ml and by qualitative nested PCR as described (8). RFLP assay was used to detect virological lamivudine resistance defined by either YIDD mutants (methionine-to-isoleucine substitution at codon 552 (M552I)) or YVDD mutants (methionine-to-valine substitution at codon 52 (M552V)) in conjunction with a leucine-to-methionine substitution at codon 528 (L528M) of the HBV polymerase gene (24, 27, 28) as previously described (25). The lower limit of detection of this method was ~500 copies viral DNA/ml serum.

Synthetic peptides

MHC class II-restricted peptides were synthesized at Epimmune on an Applied Biosystems (Foster City, CA) 430A peptide synthesizer using 9-fluorenylmethoxy carbonyl chemistry and purified to >95% purity by reverse-phase HPLC. MHC class I-restricted peptides were synthesized at Research Genetics (Huntsville, AL) at >80% purity.

Purification of HLA class II molecules

HLA class II molecules were purified from the EBV-transformed homozygous cell lines LG2 (allele DRB1*0101 (Ag DR1)); MAT (DRB1*0301 (DR3)); PREISS (DRB1*0401 (DR4w4)); KT3 (DRB1*0405 (DR4w15)); SWIG (DRB1*1101 (DR5w11)); Herluf (DRB1*1201 (DR5w12)); HO301 (DRB1*1302 (DR6w19)); PITOUT (DRB1*0701 (DR7)); OLL (DRB1*0802 (DR8w2)); HID (DRB1*0901 (DR9)); GM3107 (DRB5*0101 (DR2w2a)); TR81.9 (DRB3*0101 (DR52a)); L257.6 (DRB4*0101 (DRw53)); and from the transfected fibroblast line L466.1 (DRB1*1501 (DR2w2b)) (29). Large quantities of cells were grown in spinner cultures in RPMI 1640 with 2 mM L-glutamine (Invitrogen Life Technologies, Carlsbad, CA), 100 µg/ml streptomycin, 100 U/ml penicillin (Irvine Scientific, Santa Ana, CA), 50 µM 2-ME, and 10% heat-inactivated FCS (Irvine Scientific), and lysed for 30 min at 4°C with 50 mM Tris-HCl (pH 8.5), 1% Nonidet P-40 (Fluka Biochemika, Buchs, Switzerland), 150 mM NaCl, and 2 mM PMSF (Calbiochem, La Jolla, CA). Lysates were cleared of debris and nuclei by centrifugation at 15,000 × g for 30 min. Class II molecules were purified by affinity chromatography using the mAb LB3.1 coupled to Sepharose CL-4B beads as previously described (30, 31).

HLA-DR peptide-binding assays

Fourteen different purified human HLA-DR molecules (5–500 nM) were incubated with various unlabeled HBV peptides and 1–10 nM ¹²⁵I-radio-labeled probe peptides for 48 h. Assays were performed at pH 7.0 with the exception of that for DRB1*0301, which was performed at pH 4.5.

Table 1. MHC class II-restricted peptides

Peptide Number	HBV Protein	Amino Acid Position	Amino Acid Sequence ^a	Conservation ^b (%)
1	Nuc	50–70	PHHTALRQAILCWGELMTLA	90
2	Env	180–195	AGFFLLTRILTI PQS	80
3	Env	339–354	LVPFVQWFVGLSPTV	95
4	Pol	96–111	VGPLTVNEKRRKLKI	60
5	Pol	145–160	RHYLHTLWKAGILYK	100
6	Pol	385–400	ESRLVVDVFSQFSRGN	45
7	Pol	412–427	LQSLTNLSSNLWSL	90
8	Pol	420–435	SSNLWSLSDVSAAF	85
9	Pol	501–516	LHLYSHPIILGFRKI	80
10	Pol	618–633	KQCFRKL PVNRPIDW	45
11	Pol	664–679	KQAFTFSPITYKAFLC	60
12	Pol	694–709	LCQVFADATPTGWGL	95
13	Pol	767–782	AANWILRGTSFYVVP	70

^a Amino acid sequences of nested MHC class I binding motifs are underlined.

^b Sequence conservation among 20 full-length HBV genomes in the GenBank database, including adr, adw, ayr, and ayw isolates.

Table II. HLA-DR binding affinity

Peptide Number	HLA-DR Binding Capacity (IC ₅₀ nM) ^a												DR Bound ^b		
	DRB1*0101	DRB1*0301	DRB1*0401	DRB1*0405	DRB1*0701	DRB1*0802	DRB1*0901	DRB1*1101	DRB1*1201	DRB1*1302	DRB1*1501	DRB3*0101	DRB4*0101	DRB5*0101	% Tested
1	70	— ^c	85	—	676	196	2273	—	263	193000	9	—	—	211	7
2	1	—	8	253	8	188	58	—	6	10	217	—	—	1053	9
3	385	—	300	27	2717	74	30	—	53	1944	13	—	—	1429	7
4	8415	43	3916	1908	—	5354	—	6666	—	4461	4153	—	8121	4330	1
5	17	—	2271	1499	61	36	133	42	149	766	4	—	782	35	10
6	7372	36	208	251	6	—	—	—	—	946	1368	2525	8711	—	4
7	2	—	10	47	173	598	791	303	397	143	21	1837	4179	1067	50
8	38	62	168	17	5063	1065	7126	4923	1859	36	3089	5	7	—	7
9	248	—	77	244	—	800	1551	492	9462	—	558	—	102	560	54
10	3	—	40	34	62	872	5175	1617	—	821	4370	—	3060	1246	8
11	10	—	88	181	90	416	142	82	—	190	41	4848	322	144	6
12	7470	67	490	1203	—	—	—	—	—	2022	5009	1808	1044	—	11
13	55	—	966	1634	44	214	299	1520	802	143	386	—	6553	3279	2
															25
															8
															67

^a IC₅₀ represents the amount of peptide required for 50% inhibition of binding of the fluorescein-labeled reference peptide to each HLA-DR molecule.^b Number of DR alleles bound ≤ 1000 nM. The binding capacities < 1000 nM are indicated in bold.^c —, Indicates an IC₅₀ of > 20 μ M.

HLA-DR peptide complexes were separated from free peptide by gel filtration on TSK200 columns (TosoHaas, Montgomeryville, PA), and the fraction of bound peptide was calculated as previously described (31). In preliminary experiments, the titers of the HLA-DR preparation were determined in the presence of fixed amounts of radiolabeled peptides to determine the concentration of HLA-DR molecules necessary to bind 10–20% of the total radioactivity. Peptide inhibitors were typically tested at concentrations ranging from 1.2 ng/ml to 120 μ g/ml. All subsequent inhibition and direct binding assays were then performed using these HLA-DR concentrations.

The radiolabeled probes were HA Y307-319 for DRB1*0101; TT830-843 for DRB5*0101, DRB1*1101, DRB1*0701, DRB1*0802, and DRB1*0901; MBP Y85-100 for DRB1*1501; MT 65 KD Y3-13 with Y7 replaced with F for DRB1*0301; a nonnatural peptide with the sequence YARFQQTTLKQKT for DRB1*0401 and DRB1*0405; a nonnatural peptide with the sequence YARFQRQTLKA for DRB4*0101 (29); a naturally processed peptide of unknown origin eluted from a DRB1*1201⁺ CIR cell line with the sequence EALHQLKINPYVLS (32), an analog of TT830-843 with the sequence QYIKANAKFIGITE for DRB1*1302 (33); and integrin β_3 Y24-37 with the sequence YAWASDEALPLGSPR for DRB3*0101.

ELISPOT assay

IFN- γ ELISPOT assays were performed as previously described with duplicate cultures of 3×10^5 freshly isolated PBMC (34), 10 μ g/ml peptides, 1 μ g/ml HBV core protein (ViroGen, Watertown, MA), 50 μ g/ml tetanus toxoid (University of Massachusetts Medical School, Jamaica Plain, MA), or 1 μ g/ml PHA (Murex Biotech Limited, Dartford, U.K.) in RPMI 1640 containing 5% AB serum and 2 mM L-glutamine. Stimulation with PHA always resulted in a vigorous response with spots too numerous to count. In selected experiments, 10^5 CD4⁺ or CD8⁺ T cells, purified with anti-human CD8 Ab-conjugated magnetic microbeads (Dynal, Oslo, Norway), were cultured with 2×10^5 autologous irradiated (3000 rad) PBMCs as APCs in the presence or absence of Ags. Control experiments confirmed that the irradiated PBMC fraction did not produce any detectable IFN- γ spots. The number of specific spots (spots in the presence of Ag minus spots in the absence of Ag) was determined with a KS ELISPOT reader (Zeiss, Thornwood, NY), and represented the number of spots in the absence of Ag from the number of spots in the presence of Ag.

PBMC of 16 healthy, anti-HCV negative control persons who were negative for serum HBV DNA, HBsAg, and anti-HBc were tested with the same peptides with the following results (mean number of spot-forming cells \pm SD per 3×10^5 PBMC): peptide 1, 1.31 ± 2.4 ; NS3, peptide 2, 0.56 ± 1.2 ; peptide 3, 2.25 ± 4.2 ; peptide 4, 1 ± 1.2 ; peptide 5, 1.25 ± 2.7 ; peptide 6, 0.31 ± 1 ; peptide 7, 1.94 ± 2.8 ; peptide 8, 0.44 ± 1.1 ; peptide 9, 2.06 ± 2.8 ; peptide 10, 2.56 ± 4.1 ; peptide 11, 0.31 ± 0.8 ; and peptide 12, 0.81 ± 1.4 . A response was scored as positive if it was 1) greater than the mean response plus 2 SD in healthy, anti-HCV negative control subjects and 2) greater than 10 specific spots (spots in the presence of Ag minus spots in the absence of Ag) (35).

Cytotoxicity analysis using nested CD8⁺ T cell epitopes

HBV-specific CD8⁺ T cells were expanded from PBMC by stimulation with the nested peptides that contained HLA-A2 and HLA-A3 binding motifs (Table I). Cytotoxicity was assessed after 3 wk of culture using 25 - μ Ci-labeled, peptide-pulsed HMYC1R cells transfected with HLA-A2.1 or autologous PHA-stimulated T cell blasts as previously described (34). Spontaneous release was $< 15\%$ of maximum release in all experiments.

Statistical analysis

Fisher's exact test (two-tailed) was used to compare the frequency of immune responses in different patient groups. Student's *t* test (two-tailed) was used to compare the level of ALT and HBV DNA in different patient groups. A value of $p < 0.05$ was considered significant.

Results

Selection of highly cross-reactive HLA-DR binding peptides

Ten HBV polymerase peptides that contained either the HLA-DR supermotif (36) (peptides 5, 7, 9–11, and 13) or the DR3 motif (37) (peptides 4, 6, 8, and 12) were tested for binding to a panel of HLA DR B1, B3, B4, and B5 alleles (38). These HLA-DR molecules were selected as representative of the most common HLA variants expressed in the majority of the world's ethnic population

(38). The sequences of most selected HBV polymerase peptides were conserved in >80% of the 20 most common HBV strains including adr, adw, ayr, and ayw isolates (Table I). In addition, several peptides that were conserved in only ~50% of isolates, but highly conserved in their DR binding region, were included in the analysis. For comparison, we also studied one HBV nucleocapsid (39) and two HBV envelope peptides (40, 41) (Table I) and the HBcore protein (see Fig. 4).

As shown in Table II, all HBV peptides bound with an IC_{50} of ≤ 1000 nM to at least one HLA-DR molecule. The IC_{50} value represents the amount of peptide required for 50% inhibition of binding of a fluorescein-labeled reference peptide. An IC_{50} value of 1000 nM has previously been shown to represent the affinity threshold associated with immunogenicity (38, 42). Moreover, 8 of the 10 polymerase peptides bound at least four of these frequent DR molecules. This degree of cross-reactivity is not dissimilar from the previously identified control peptides (39–41), and suggested that these peptides should be recognized by subjects with diverse HLA haplotypes. Based on the worldwide prevalence of the HLA-DR Ags, this panel of peptides should cover 98.2% of the average population.

Immunogenicity of HBV polymerase peptides as assessed by IFN- γ ELISPOT analysis

To determine whether these highly cross-reactive and conserved HLA-DR binding peptides were recognized by T cells of patients with past or present HBV infection, IFN- γ ELISPOT responses were evaluated with PBMC from 1 acutely HBV-infected patient, 17 recovered patients, and 48 persistently HBV-infected patients (Table III). Sixteen blood donors were tested as controls. The cut-off of positivity was set at ≥ 10 IFN- γ spots/300,000 PBMC (see *Materials and Methods*). All 10 HBV polymerase peptides were recognized by PBMC of at least one patient, and 29 of 66 patients (44%) responded to at least one of the analyzed HBV polymerase peptides (Fig. 1). Peptides 4 and 8 were the only peptides that were exclusively recognized by the patient with strongest immune response and acute hepatitis B, and one of these peptides (peptide 4) had a very low HLA-DR binding affinity (Table II). By contrast, peptides 7 and 11, associated with significant binding to 77 and 92% of the HLA-DR molecules, respectively, were recognized by 8 and 13 patients. These data demonstrate that conserved HBV polymerase peptides selected on the basis of a high binding affinity to several HLA DR molecules are antigenic in the context of multiple DR alleles, and thus allow coverage of a broad and ethnically diverse patient population.

In the total patient population, 61 responses were observed against HBV polymerase peptides. The patients' HLA haplotypes

were known for 56 of these 61 cases. When the observed responses were evaluated in the context of the patients' HLA haplotypes, 55% (31 of 56) of all responses occurred in the presence of one of the DR alleles with high peptide-binding affinity ($IC_{50} < 1000$ nM; Table II). In an additional 20% (11 of 56) of all responses, the patients expressed DR molecules at least weakly binding (1000 nM $< IC_{50} < 20$ μ M) the peptide. In the remaining cases, patients did not express any DR alleles known to bind the peptide ($IC_{50} < 20$ μ M; Table II), suggesting that some of the responses might be restricted by other alleles not analyzed in the current study.

The magnitude of the HBV-specific T cell responses was assessed by the frequency of HBV nucleocapsid, envelope and polymerase peptide-specific, IFN- γ -producing T cells in the PBMC population (Fig. 2). The broadest T cell response was noted during acute, self-limited hepatitis B with responses to all 13 HBV epitopes. In contrast, none of the other patients responded to more than five epitopes, and the overall response of patients with chronic hepatitis B was weak. As previously reported for other HBV epitopes and proteins (35, 43, 44), the frequency of HBV epitope-specific cells was low, i.e., between ~10 and 50 peptide-specific cells per 300,000 PBMC. Tetanus toxoid-specific responses were tested as a positive control, and the frequency of responses did not differ significantly among patient subgroups (Fig. 2C).

T cell subsets responding to HBV polymerase peptides

To determine the T cell subset that responded to the HBV polymerase peptides, PBMC were separated into CD4⁺ and CD8⁺ subpopulations using Ab-coated magnetic beads. Each cell subpopulation was then separately tested in the presence of irradiated, autologous APCs and the respective peptides. Fig. 3A demonstrates in a representative experiment that most IFN- γ spots in the ELISPOT assay segregated with the CD4⁺ T cell subfraction. However, in the case of peptide 9, ~25% of the total number of IFN- γ spots could still be attributed to CD8⁺ T cells (Fig. 3A). A closer analysis of the peptide sequences revealed that peptide 9 as well as four other polymerase peptides contained nested HLA class I-binding motifs (Table I). In fact, nested sequences within peptides 5, 6, and 11 have previously been described as CD8⁺ T cell epitopes (45). Thus, in natural HBV infection, these shorter peptides may be endogenously processed from polypeptide precursors, loaded onto HLA class I molecules in HBV-infected cells, transported to the cell surface, and stimulate responses of CD8⁺ T cells.

To test this hypothesis, we chose the nested sequences HLY-SHPIL in peptide 9 and QAFTFSPTYK in peptide 11, which contained the HLA-A2 and HLA-A3 binding motifs, respectively, and tested them for CTL recognition using a microwell peptide stimulation technique to expand low-frequency HBV-specific CTL

Table III. Baseline characteristics of the patients studied

Group	Clinical Diagnosis	No. of Patients	Sex M/F ^a	Race C/A/AA/M ^b	Age (yr) Mean \pm SD	ALT (IU/L) Mean \pm SD	HBsAg ⁺ n (%)	HBV DNA (copies/ml)
1	Acute HBV patient	1	1/0	1/0/0/0	27	641	1 (100)	1.5×10^4
2	Recovered patients	17	15/2	14/1/2/0	52 \pm 9	55 \pm 12	0 (0)	0 ^c
3	Chronic hepatitis B, no therapy							
3a	HBsAg positive	7	6/1	4/2/0/1	45 \pm 21	97 \pm 58	7 (100)	$3.2 \times 10^8 \pm 2.8 \times 10^8$
3b	HBsAg negative	7	4/3	3/4/0/0	49 \pm 9	87 \pm 87	7 (100)	$2.5 \times 10^7 \pm 5.2 \times 10^7$
4	Chronic hepatitis B, on therapy ^d							
4a	HBsAg positive	20	18/2	13/6/1/0	46 \pm 3	84 \pm 82	20 (100)	$2.0 \times 10^8 \pm 2.7 \times 10^8$
4b	HBsAg negative	14	12/2	5/6/1/2	49 \pm 13	41 \pm 21	14 (100)	$3.4 \times 10^5 \pm 9.5 \times 10^5$

^a M, Male; F, female.

^b C, Caucasian; A, Asian; AA, African-American; M, Mediterranean.

^c Negative by branched DNA analysis and nested PCR.

^d Lamivudine therapy (100 mg orally per day).

Patient Group	Pt. No.	IFN- γ ELISPOT Responses													Response Frequency per Peptide		
		Nucleocapsid and Envelope Peptides			Polymerase Peptides												
		1	2	3	4	5	6	7	8	9	10	11	12	13	[n]	[%]	
1 Acute, self-limited 2 Recovered	1														13	100%	
	2														5	38%	
	3														2	15%	
	4														1	8%	
	5														5	38%	
	7														5	38%	
	9														3	23%	
	10														3	23%	
	11														1	8%	
	12														3	23%	
	13														3	23%	
	14														1	8%	
	15														1	8%	
	16														1	8%	
	17														1	8%	
	18														1	8%	
	3a Chronic, untreated HBeAg positive	19														1	8%
		22														1	8%
23															2	15%	
3b Chronic, untreated HBeAg negative	26														1	8%	
	27														4	31%	
	29														2	15%	
	32														1	8%	
4a Chronic, treated HBeAg positive	38														3	23%	
	40														3	23%	
	41														5	38%	
	42														1	8%	
	45														3	23%	
	48														5	38%	
4b Chronic, treated HBeAg negative	53														3	23%	
	54														3	23%	
	58														2	15%	
	60														1	8%	
	62														2	15%	
	63														1	8%	
	Response Frequency [n] Per Patient		8	9	17	1	2	13	8	1	4	9	13	4	6		
		9%	14%	26%	2%	3%	20%	12%	2%	6%	14%	20%	6%	9%			

FIGURE 1. HBV nucleocapsid, envelope, and polymerase peptide-specific T cell responses. Only patients who recognized at least one of the 13 tested peptides in the IFN- γ ELISPOT assay are shown. A positive IFN- γ response was defined as >10 spot-forming cells per 300,000 PBMC, which is greater than the mean plus 2 SD of the baseline response detected in 16 healthy uninfected controls. Shaded boxes indicate the presence of a significant IFN- γ T cell response to HBV nucleocapsid and envelope peptides; black boxes indicate the presence of a significant IFN- γ T cell response to HBV polymerase peptides. Peptide sequences are described in Table I, and patient groups in Table III.

precursors from PBMC. Overall, the HLA-A2 restricted epitope in peptide 9 and the HLA-A3-restricted epitope in peptide 11 expanded HBV-specific CD8⁺ T cells of several patients with the corresponding HLA class I haplotype (Fig. 3B). These results demonstrate that several of the newly identified CD4⁺ T cell epitopes contained shorter sequences that were recognized by CD8⁺ T cells when presented as the minimal optimal epitope in the context of the matching HLA class I molecule.


Cross-sectional analysis of HBV polymerase-specific T cell responses in patient groups with different outcome of HBV infection

Because HBV polymerase is an essential enzyme for HBV replication, expressed early in the HBV life cycle and also found inside the secreted HBV particle, we reasoned that the frequency of HBV-specific T cell responses might correlate with different stages of HBV infection. As previously shown for CD8⁺ T cell responses against polymerase (2, 3), there was a significant difference in the frequency of CD4⁺ T cell responses between the different patient subgroups. Whereas the patient with acute, self-limited hepatitis B recognized all epitopes, and 11 of 17 (65%) recovered patients (group 2) tested positive for at least one HBV polymerase peptide, only 16 of 48 (33%) of patients with chronic hepatitis B (groups 3 and 4) tested positive ($p = 0.043$; Fig. 1). When immune responses against any peptide, i.e., including the nucleocapsid and envelope peptides, were considered, this difference between

acutely infected and recovered patients (16 of 18 (89%) responding) and chronically infected patients (19 of 48 (40%) responding) was even more significant ($p < 0.0001$; Fig. 1).

Among patients with chronic hepatitis B, further analysis was performed to assess the role of lamivudine therapy on cellular immune responses to HBV polymerase peptides. Cellular immune responses to HBV polymerase were detected in 4 of 14 untreated (29%) and in 12 of 34 (35%) treated patients. Within the lamivudine-treated group, the duration of treatment at the time of testing appeared to correlate with reactivity. Four of 5 (80%) patients treated for <1 year responded to the polymerase epitopes as compared with 3 of 7 (43%) patients treated for 1–3 years, 5 of 22 (24%) patients treated for >3 years, and 4 of 14 (29%) patients who were not treated (Fig. 4A). These differences in the polymerase-specific responses between the subgroups were statistically significant ($p = 0.029$ for <1 -year treatment vs >3 -year treatment; $p = 0.041$ for <1 -year treatment vs >1 -year treatment) (Fig. 4A) and correlated with differences in viral levels among patient subgroups. For example, the highest frequency of HBV polymerase-specific T cell responses was observed in patients who had received lamivudine therapy for <1 year and displayed the lowest HBV DNA levels. In contrast, HBV polymerase-specific T cell responses were lower in patients who had been treated for 1–3 and for 3–5 years, respectively, and these patient subgroups were characterized by 100- to 1000-fold higher HBV DNA levels due to development of lamivudine-resistant HBV mutants (Fig. 4A).

FIGURE 2. Vigor of HBV nucleocapsid, envelope, and polymerase peptide-specific T cell responses. *A* and *B*, Direct ex vivo IFN- γ ELISPOT analysis of peripheral blood T cell responses to HBV nucleocapsid and envelope peptides (*A*) and HBV polymerase peptides (*B*). Only significant IFN- γ responses greater than the mean plus 2 SD (indicated by horizontal line) of the baseline response detected in 16 healthy uninfected controls and >10 specific spots/300,000 PBMC are shown. The asterisk indicates 102 spots. Peptide sequences are described in Table I, patient groups (■, group 1; □, group 2; ▨, group 3; ▩, group 4) are defined in Table III. *C*, Tetanus-specific responses. Responses of >100 spots were too numerous to count with the automated ELISPOT reader.



Overall, the frequency of HBV polymerase-specific responses was higher in patients who had successfully cleared HBeAg (Fig. 4*B*) than in patients who remained HBeAg positive under lamivudine therapy (*B*). When HBeAg⁺ and HBeAg⁻ patients were analyzed separately, HBV polymerase-specific T cell response decreased in both subgroups under prolonged lamivudine therapy. Consistent with the findings for all treated patients (group 4), this reduced T cell responsiveness correlated with increased HBV DNA levels and emergence of lamivudine resistance mutants, but did not reach statistical significance due to the small number of patients in each subgroup (Fig. 4*B*). Finally, a similar trend toward decreased HBV-specific immune responsiveness under prolonged lamivudine therapy was observed for HBcore-specific responses (Fig. 4*C*), whereas responses to the control Ag tetanus toxoid were comparable for all patients (Fig. 4*D*).

Collectively, the data suggest that antiviral therapy restores HBV polymerase-specific immune responses transiently, i.e., during the first year of therapy, and that cellular immune responses gradually attenuate with emergence of lamivudine-resistant HBV mutants and increase of viral levels under more prolonged therapy.

Discussion

The current study uses a combined immunochemical and cellular immunology approach to identify and characterize 10 highly conserved and immunogenic CD4⁺ T cell epitopes in the HBV polymerase protein. The newly identified CD4⁺ T cell epitopes within HBV polymerase are relevant for several reasons. First, all epitopes displayed significant binding affinity to multiple HLA-DR molecules. Therefore, the data suggest that the majority of the world's population, independent of ethnicity, should be able to respond to these epitopes. Second, CD4⁺ T cell epitopes within HBV polymerase offer significant advantages for the development of epitope-based diagnostics and vaccines, because of the high

degree of sequence conservation of HBV polymerase and its essential role in the viral life cycle, which reduces the possibility of viral escape by mutations in T cell epitopes. In addition, HBV polymerase represents the largest HBV protein and has already been shown to be highly immunogenic at the CD8⁺ T cell level in patients with acute, self-limited hepatitis B (2, 3). Third, the newly




FIGURE 3. Identification of nested MHC class I-restricted epitopes. *A*, PBMC were sorted into CD4⁺ and CD8⁺ T cell subpopulations using Ab-coated magnetic beads and stimulated separately with peptide-pulsed autologous APCs. The number of peptide-specific, IFN- γ -producing cells as determined by ELISPOT analysis is indicated and greater for the CD4⁺ T cell subpopulation than for the CD8⁺ T cell subpopulation. A representative experiment is shown. *B*, Cytotoxic CD8⁺ T cells specific for MHC class I-restricted peptides that are nested within the HBV-polymerase CD4⁺ T cell epitopes. T cell lines were generated from PBMC of the HLA-A2-positive patients 12, 29, and 39, and the HLA-A3-positive patient 17, respectively, by stimulation with the nested 9- and 10-mer peptides located within peptide 9 (●) or 11 (■) (see Table I). Expanded T cell lines were then tested for specific cytotoxicity against the nested peptides in a standard ⁵¹Cr release assay at the indicated E:T ratios.




FIGURE 4. Prevalence of HBV polymerase (A and B), HBcore (C), and tetanus toxoid-specific (D) T cell responses in recovered patients (group 2), untreated patients with chronic hepatitis B (group 3), and patients with chronic hepatitis B during lamivudine therapy (group 4). Patients with chronic hepatitis B are separated into those with HBeAg or without HBeAg in B.

identified HBV polymerase epitopes were able to induce IFN- γ production by HBV-specific T cells, an effector function associated with in vivo suppression of HBV replication (7, 46). Fourth, the observation that several of these CD4⁺ T cell epitopes contain nested MHC class I-restricted epitopes that stimulate IFN- γ responses and cytotoxicity by CD8⁺ T cells is an additional attractive feature for a multiepitope vaccine. The well-characterized, immunodominant HBe₁₈₋₂₇ CD8⁺ T cell epitope, for example, does also overlap with an MHC class II-restricted CD4⁺ T cell epitope (47), and it has been described that CD4⁺ T cells are indispensable for the maintenance of functional CD8⁺ T cells that control chronic viral infections (48). This CD4⁺ T cell help may consist of direct help for CD8⁺ T cells via production of cytokines (49) or of indirect help for professional APCs via CD40/CD40L-mediated activation (50). Moreover, recent studies have demonstrated the need to include potent CD4⁺ T cell epitopes to restore an altered Th response and to overcome CD8⁺ T cell tolerance in chronic HBV infection (14). Because covalent linkage of CD4⁺ and CD8⁺ T cell epitopes on the same peptide vaccine construct is important for the induction of Ag-specific responses (51), the natural occurrence of nested CD8⁺ and CD4⁺ T cell epitopes within the polymerase Ag might provide an elegant and promising approach to fulfill these requirements.

Finally, the presence of HBV polymerase-specific, CD4⁺ T cell responses correlated with different outcomes of HBV infection. As demonstrated for CD8⁺ T cell responses (2), circulating HBV polymerase-specific CD4⁺ T cell responses were more frequent in acute self-limited hepatitis B and after recovery than in chronic hepatitis B. Moreover, even in the persistently infected, HBsAg⁺ patient subgroup, HBV polymerase-specific T cell responses were associated with a partial response, i.e., with loss of HBeAg (Fig.

4B) and reduction of HBV DNA levels (A). Although the recognition of endogenously processed HBV polymerase could not be tested due to a lack of recombinant and purified polymerase protein, these observations and the absence of those responses in healthy, uninfected blood donors indicated that all epitopes were processed in vivo and that HBV polymerase-specific CD4⁺ T cells were primed in vivo.

In the second part of this study, the newly identified CD4⁺ T cell epitopes were then used to analyze the HBV-specific immune response during antiviral therapy. Recent studies have suggested that suppression of HBV replication by lamivudine rapidly restores HBV-specific cellular immune responsiveness to the same level as in recovered persons (20). In that study, both CD4⁺ (20) and CD8⁺ (21) T cells specific for HBV remained detectable in the blood for up to 5 mo during lamivudine therapy. However, the increased T cell reactivity was not associated with an increase in loss of HBeAg or HBsAg (20), the serological hallmark of recovery from hepatitis B. Also, a second study of patients receiving IFN- α alone and in combination with lamivudine reported no restoration or de novo induction of HBV core-specific T cell proliferation (22). Because HBV polymerase is expressed early in the HBV life cycle and also found inside the secreted HBV particle, we reasoned that the frequency of HBV polymerase-specific T cells might correlate more closely with viral levels and the outcome of lamivudine therapy.

Using the newly identified HBV polymerase peptides, the current study indicated that antiviral therapy of persistently infected patients appeared to increase the frequency of HBV-specific CD4⁺ T cell responses during the first year of treatment. This result indicates that HBV-specific T cells are not completely depleted or anergized during chronic hepatitis B because their frequency in the

peripheral blood can significantly increase upon therapeutic reduction of HBV levels. The reconstitution of HBV-specific T cells in the periphery may result from generation of new precursors in the regional lymph nodes on the one hand (52) and from decreased sequestration of effector T cells to the liver.

With longer periods of antiviral therapy, however, the frequency of HBV-specific CD4⁺ T cell responses decreased and, after 3 years of therapy, responses were no more frequent than in untreated patients. Decreased responsiveness of HBV-specific T cells correlated with the emergence of lamivudine-resistant HBV mutants and a concomitant increase of HBV DNA levels. Whether the lack of HBV-specific responses predated the development of viral resistance and may have contributed to its development, is an interesting question, that could not be answered due to the cross-sectional nature of this study and warrants future, prospective studies. Overall, the results suggest that therapeutic vaccination designed to boost HBV-specific cellular immunity might be most efficient during the first year of antiviral therapy. The degenerate HLA-DR binding and immunogenicity of the HBV polymerase epitopes as well as the inclusion of nested CD8⁺ T cell epitopes make these epitopes valuable components of a vaccine that is designed to cover a large and ethnically diverse patient population.

Acknowledgments

We thank Ms. Yoon Park for invaluable help with sample collection, and all of the patients who donated blood samples for this study.

References

- Chisari, F. V., and C. Ferrari. 1995. Hepatitis B virus immunopathogenesis. In *Annual Review of Immunology*. W. E. Paul, ed. Annual Reviews, Inc., Palo Alto, CA, p. 29.
- Rehermann, B., P. Fowler, J. Sidney, P. Person, A. Redeker, M. Brown, B. Moss, A. Sette, and F. V. Chisari. 1995. The cytotoxic T lymphocyte response to multiple hepatitis B virus polymerase epitopes during and after acute viral hepatitis. *J. Exp. Med.* 181:1047.
- Maini, M. K., C. Boni, G. S. Ogg, A. S. King, S. Reingnat, C. K. Lee, J. R. Larubia, G. J. Webster, A. J. McMichael, C. Ferrari, et al. 1999. Direct ex vivo analysis of hepatitis B virus-specific CD8⁺ T cells associated with the control of infection. *Gastroenterology* 117:1386.
- Guidotti, L. G., and F. V. Chisari. 2001. Nontoxic control of viral infections by the innate and adaptive immune response. *Annu. Rev. Immunol.* 19:65.
- Kakimi, K., T. E. Lane, F. V. Chisari, and L. G. Guidotti. 2001. Cutting edge: inhibition of hepatitis B virus replication by activated NK T cells does not require inflammatory cell recruitment to the liver. *J. Immunol.* 167:6701.
- Moriyama, T., S. Guilhot, K. Klopchin, B. Moss, C. A. Pinkert, R. D. Palmiter, R. L. Brinster, O. Kanagawa, and F. V. Chisari. 1990. Immunobiology and pathogenesis of hepatocellular injury in hepatitis B virus transgenic mice. *Science* 248:361.
- Guidotti, L. G., R. Rochford, J. Chung, M. Shapiro, R. Purcell, and F. V. Chisari. 1999. Viral clearance without destruction of infected cells during acute HBV infection. *Science* 284:825.
- Rehermann, B., D. Lau, J. Hoofnagle, and F. V. Chisari. 1996. Cytotoxic T lymphocyte responsiveness after resolution of chronic hepatitis B virus infection. *J. Clin. Invest.* 97:1655.
- Ferrari, C., A. Penna, A. Bertoletti, A. Valli, A. D. Antoni, T. Giuberti, and A. Cavalli. 1990. Cellular immune response to hepatitis B virus-encoded antigens in acute and chronic hepatitis B virus infection. *J. Immunol.* 145:3442.
- Missale, G., A. Redeker, J. Person, P. Fowler, S. Guilhot, H. J. Schlicht, C. Ferrari, and F. V. Chisari. 1993. HLA-A31 and A68 restricted cytotoxic T cell responses to a single hepatitis B virus nucleocapsid epitope during acute viral hepatitis. *J. Exp. Med.* 177:751.
- Nayersina, R., P. Fowler, S. Guilhot, G. Missale, A. Cerny, H.-J. Schlicht, A. Vitiello, R. Chesnut, J. L. Person, A. G. Redeker, and F. V. Chisari. 1993. HLA A2 restricted cytotoxic T lymphocyte responses to multiple hepatitis B surface antigen epitopes during hepatitis B virus infection. *J. Immunol.* 150:4659.
- Tsai, S. L., M. Y. Chen, M. Y. Lai, P. M. Yang, J. L. Seng, J. H. Huang, L. H. Hwang, T. H. Chang, and D. S. Chen. 1992. Acute exacerbations of chronic type B hepatitis are accompanied by increased T cell responses to hepatitis B core and e antigens. *J. Clin. Invest.* 89:87.
- Vitiello, A., G. Ishioka, H. M. Grey, R. Rose, P. Farness, R. LaFond, L. Yuan, F. V. Chisari, J. Furze, R. Bartholomew, and R. W. Chesnut. 1995. Development of a lipopeptide-based therapeutic vaccine to treat chronic HBV infection. 1. Induction of a primary cytotoxic T lymphocyte response in humans. *J. Clin. Invest.* 95:341.
- Livingston, B. D., J. Alexander, C. Crimi, C. Oseroff, E. Celis, K. Daly, L. G. Guidotti, F. V. Chisari, J. Fikes, R. W. Chesnut, and A. Sette. 1999. Altered helper T lymphocyte function associated with chronic hepatitis B virus infection and its role in response to therapeutic vaccination in humans. *J. Immunol.* 162:3088.
- Livingston, B. D., C. Crimi, H. Grey, G. Ishioka, F. V. Chisari, J. Fikes, R. W. Chesnut, and A. Sette. 1997. The hepatitis B virus-specific CTL responses induced in humans by lipopeptide vaccination are comparable to those elicited by acute viral infection. *J. Immunol.* 159:1383.
- Heathcote, J., J. McHutchison, S. Lee, M. Tong, K. Benner, G. Minuk, T. Wright, J. Fikes, B. Livingston, A. Sette, and R. Chesnut. 1999. A pilot study of the CY-1899 T-cell vaccine in subjects chronically infected with hepatitis B virus: The CY1899 T Cell Vaccine Study Group. *Hepatology* 30:531.
- Ridge, J. P., F. Di Rosa, and P. Matzinger. 1998. A conditioned dendritic cell can be a temporal bridge between a CD4⁺ T-helper and a T-killer cell. *Nature* 393:474.
- Sigal, L. J., S. Crotty, R. Andino, and K. L. Rock. 1999. Cytotoxic T-cell immunity to virus-infected non-haematopoietic cells requires presentation of exogenous antigen. *Nature* 398:77.
- Kakimi, K., M. Isogawa, J. Chung, A. Sette, and F. V. Chisari. 2002. Immunogenicity and tolerogenicity of hepatitis B virus structural and nonstructural proteins: implications for immunotherapy of persistent viral infections. *J. Virol.* 76:8609.
- Boni, C., A. Bertoletti, A. Penna, A. Cavalli, M. Pilli, S. Urbani, P. Scognamiglio, R. Boehme, R. Panebianco, F. Fiaccadori, and C. Ferrari. 1998. Lamivudine treatment can restore T cell responsiveness in chronic hepatitis B. *J. Clin. Invest.* 102:968.
- Boni, C., A. Penna, G. S. Ogg, A. Bertoletti, M. Pilli, C. Cavallo, A. Cavalli, S. Urbani, R. Boehme, R. Panebianco, et al. 2001. Lamivudine treatment can overcome cytotoxic T-cell hyporesponsiveness in chronic hepatitis B: new perspectives for immune therapy. *Hepatology* 33:963.
- Marinos, G., N. V. Naoumov, and R. Williams. 1996. Impact of complete inhibition of viral replication on the cellular immune response in chronic hepatitis B virus infection. *Hepatology* 24:991.
- Leung, N. W., C. L. Lai, T. T. Chang, R. Guan, C. M. Lee, K. Y. Ng, S. G. Lim, P. C. Wu, J. C. Dent, S. Edmundson, et al. 2001. Extended lamivudine treatment in patients with chronic hepatitis B enhances hepatitis B e antigen seroconversion rates: results after 3 years of therapy. *Hepatology* 33:1527.
- Lai, C. L., R. N. Chien, N. W. Leung, T. T. Chang, R. Guan, D. I. Tai, K. Y. Ng, P. C. Wu, J. C. Dent, J. Barber, et al. 1998. A one-year trial of lamivudine for chronic hepatitis B: Asia Hepatitis Lamivudine Study Group. *N. Engl. J. Med.* 339:61.
- Lau, D. T., M. F. Khokhar, E. Doo, M. G. Ghany, D. Herion, Y. Park, D. E. Kleiner, P. Schmid, L. D. Condey, J. Gauthier, et al. 2000. Long-term therapy of chronic hepatitis B with lamivudine. *Hepatology* 32:828.
- Urdea, M. 1993. Synthesis and characterization of branched DNA (bDNA) for the direct and quantitative detection of CMV, HBV, HCV and HIV. *Clin. Chem.* 39:725.
- Bartholomew, M. M., R. W. Jansen, L. J. Jeffers, K. R. Reddy, L. C. Johnson, H. Bunzendahl, L. D. Condey, A. G. Tzakis, E. R. Schiff, and N. A. Brown. 1997. Hepatitis-B-virus resistance to lamivudine given for recurrent infection after orthotopic liver transplantation. *Lancet* 349:20.
- Allen, M. I., J. Gauthier, M. DesLauriers, E. J. Bourne, K. M. Carrick, F. Baldanti, L. L. Ross, M. W. Lutz, and L. D. Condey. 1999. Two sensitive PCR-based methods for detection of hepatitis B virus variants associated with reduced susceptibility to lamivudine. *J. Clin. Microbiol.* 37:3338.
- Alexander, J., J. Sidney, S. Southwood, J. Ruppert, C. Oseroff, A. Maewal, K. Snoke, H. M. Serra, R. T. Kubo, A. Sette, et al. 1994. Development of high potency universal DR-restricted helper epitopes by modification of high affinity DR-blocking peptides. *Immunity* 1:751.
- Gorga, J. C., V. Horejsi, D. R. Johnson, R. Raghupathy, and J. L. Strominger. 1987. Purification and characterization of class II histocompatibility antigens from a homozygous human B cell line. *J. Biol. Chem.* 262:16087.
- Sette, A., S. Buus, S. Colon, C. Miles, and H. M. Grey. 1989. Structural analysis of peptides capable of binding to more than one Ia antigen. *J. Immunol.* 142:35.
- Falk, K., O. Rotzschke, S. Stevanovic, G. Jung, and H. G. Rammensee. 1994. Pool sequencing of natural HLA-DR, DQ, and DP ligands reveals detailed peptide motifs, constraints of processing, and general rules. *Immunogenetics* 39:230.
- Boitel, B., U. Blank, D. Mege, G. Corradin, J. Sidney, A. Sette, and O. Acuto. 1995. Strong similarities in antigen fine specificity among DRB1*1302-restricted tetanus toxin tt830-843-specific TCRs in spite of highly heterogeneous CDR3. *J. Immunol.* 154:3245.
- Takaki, A., M. Wiese, G. Maertens, E. Depla, U. Seifert, A. Liebetrau, J. L. Miller, M. P. Manns, and B. Rehermann. 2000. Cellular immune responses persist, humoral responses decrease two decades after recovery from a single source outbreak of hepatitis C. *Nat. Med.* 6:578.
- Jung, M. C., B. Hartmann, J. T. Gerlach, H. Diepolder, R. Gruber, W. Schraut, N. Gruner, R. Zachoval, R. Hoffmann, T. Santantonio, et al. 1999. Virus-specific lymphokine production differs quantitatively but not qualitatively in acute and chronic hepatitis B infection. *Virology* 261:165.
- O'Sullivan, D., T. Arhenius, J. Sidney, M. F. Del Guercio, M. Albertson, M. Wall, C. Oseroff, S. Southwood, S. M. Colon, F. C. A. Gaeta, and A. Sette. 1991. On the interaction of promiscuous antigenic peptides with different DR alleles: identification of common structural motifs. *J. Immunol.* 147:2663.
- Southwood, S., J. Sidney, A. Kondo, M. F. del Guercio, E. Appella, S. Hoffman, R. T. Kubo, R. W. Chesnut, H. M. Grey, and A. Sette. 1998. Several common HLA-DR types share largely overlapping peptide binding repertoires. *J. Immunol.* 160:3363.
- Wilson, C. C., B. Palmer, S. Southwood, J. Sidney, Y. Higashimoto, E. Appella, R. Chesnut, A. Sette, and B. D. Livingston. 2001. Identification and antigenicity

- of broadly cross-reactive and conserved human immunodeficiency virus type 1-derived helper T-lymphocyte epitopes. *J. Virol.* 75:4195.
39. Penna, A., M. Artini, A. Cavalli, M. Levrero, A. Bertoletti, M. Pilli, F. V. Chisari, B. Rehmann, G. Del Prete, F. Fiaccadori, and C. Ferrari. 1996. Long lasting memory T cell responses following self-limited acute hepatitis B. *J. Clin. Invest.* 98:1185.
40. Celis, E., D. Ou, and L. Otvos. 1988. Recognition of hepatitis B surface antigen by human T lymphocytes: proliferative and cytotoxic responses to a major antigenic determinant defined by synthetic peptides. *J. Immunol.* 140:1808.
41. Barnaba, V., A. Franco, M. Paroli, R. Benvenuto, G. De Petrillo, V. L. Burgio, and I. Santilio. 1994. Selective expansion of cytotoxic T lymphocytes with a CD4⁺CD56⁺ surface phenotype and a T helper type 1 profile of cytokine secretion in the liver of patients chronically infected with hepatitis B virus. *J. Immunol.* 153:5324.
42. Sette, A., A. Vitiello, B. Rehmann, P. Fowler, R. Nayersina, C. Oseroff, L. Yuan, J. Ruppert, J. Sidney, M.-F. del Guercio, et al. 1994. The relationship between class I binding affinity and immunogenicity of potential cytotoxic T cell epitopes. *J. Immunol.* 153:5586.
43. Hyodo, N., M. Tajimi, T. Ugajin, I. Nakamura, and M. Imawari. 2003. Frequencies of IFN- γ and interleukin-10 secreting cells in peripheral blood mononuclear cells and liver infiltrating lymphocytes in chronic hepatitis B virus infection. *Hepatol. Res.* 27:109.
44. Szkaradkiewicz, A., A. Jopek, J. Wysocki, M. Grzymislawski, I. Malecka, and A. Wozniak. 2003. HBcAg-specific cytokine production by CD4 T lymphocytes of children with acute and chronic hepatitis B. *Virus Res.* 97:127.
45. Bertoni, R., J. Sidney, P. Fowler, R. W. Chesnut, F. V. Chisari, and A. Sette. 1997. Human histocompatibility leukocyte antigen-binding supermotifs predict broadly cross-reactive cytotoxic T lymphocyte responses in patients with acute hepatitis. *J. Clin. Invest.* 100:503.
46. Guidotti, L. G., S. Guilhot, and F. V. Chisari. 1994. Interleukin 2 and interferon α/β downregulate hepatitis B virus gene expression in vivo by tumor necrosis factor dependent and independent pathways. *J. Virol.* 68:1265.
47. Bertoletti, A., S. Southwood, R. Chesnut, A. Sette, M. Falco, G. B. Ferrara, A. Penna, C. Boni, F. Fiaccadori, and C. Ferrari. 1997. Molecular features of the hepatitis B virus nucleocapsid T-cell epitope 18–27: interaction with HLA and T-cell receptor. *Hepatology* 26:1027.
48. Zajac, A. J., J. N. Blattman, K. Murali-Krishna, D. J. Sourdive, M. Suresh, J. D. Altman, and R. Ahmed. 1998. Viral immune evasion due to persistence of activated T cells without effector function. *J. Exp. Med.* 188:2205.
49. Lu, Z., L. Yuan, X. Zhou, E. Sotomayor, H. I. Levitsky, and D. M. Pardoll. 2000. CD40-independent pathways of T cell help for priming of CD8⁺ cytotoxic T lymphocytes. *J. Exp. Med.* 191:541.
50. Bennett, S. R., F. R. Carbone, F. Karamalis, R. A. Flavell, J. F. Miller, and W. R. Heath. 1998. Help for cytotoxic-T-cell responses is mediated by CD40 signalling. *Nature* 393:478.
51. Shirai, M., C. D. Pendleton, J. Ahlers, T. Takeshita, M. Newman, and J. A. Berzofsky. 1994. Helper-cytotoxic T lymphocyte (CTL) determinant linkage required for priming of anti-HIV CD8⁺ CTL in vivo with peptide vaccine constructs. *J. Immunol.* 152:549.
52. Malacarne, F., G. J. Webster, S. Reignat, J. Gotto, S. Behboudi, A. K. Burroughs, G. M. Dusheiko, R. Williams, and A. Bertoletti. 2003. Tracking the source of the hepatitis B virus-specific CD8 T cells during lamivudine treatment. *J. Infect. Dis.* 187:679.